

Stroke Guideline

History

- Previous CVA, TIAs
- Previous cardiac / vascular surgery
- Associated diseases: diabetes, hypertension, CAD
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma

Signs and Symptoms

- Altered mental status
- Weakness / Paralysis
- Acute focal neuro deficit
- Blindness or other sensory loss
- Aphasia / Dysarthria
- Syncope
- Vertigo / Dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension / hypotension


Differential

- See Altered Mental Status
- TIA (Transient ischemic attack)
- Seizure
- Todd's Paralysis
- Hypoglycemia
- Stroke
- Tumor
- Trauma
- Dialysis / Renal Failure

SEVERITY-BASED STROKE TRIAGE ALGORITHM FOR EMS



Together to End Stroke™



Adult Medical Guidelines

CPSS=Gincinnati Pre-hospital Stroke Scale, LAPSS=LA Prehospital Stroke Scale LVO=Large Vessel Occlusion, LKW=Last Known Well
EVT=Endovascular therapy, ASRH=Acute Stroke Ready Hospital, PSC=Primary Stroke Center, CSC=Comprehensive Stroke Center

Stroke Guideline

Cincinnati Pre-hospital Stroke Scale

1. FACIAL DROOP: Have patient show teeth or smile.



Normal:
both sides
of the face
move equally



Abnormal:
one side of
face does not
move as well
as the other
side

2. ARM DRIFT: Patient closes eyes & holds both arms out for 10 sec.



Normal:
both arms
move the
same or both
arms do not
move at all



Abnormal:
one arm does
not move or
drifts down
compared to
the other

3. ABNORMAL SPEECH: Have the patient say "you can't teach an old dog new tricks."

Normal: patient uses correct words with no slurring

Abnormal: patient slurs words, uses the wrong words, or is unable to speak

INTERPRETATION: If any 1 of these 3 signs is abnormal, the probability of a stroke is 72%.

| ITEM | Instruction | Result | Score |
|--|--|---|-------|
| Facial Palsy | Ask patient to show their teeth (smile) | Absent (symmetrical movement) | 0 |
| | | Mild (slight asymmetrical) | 1 |
| | | Moderate to Severe (completely asymmetrical) | 2 |
| Arm Motor Function | Extending the arm of the patient 90° (if sitting) or 45° (if supine) | Normal to Mild (limb upheld more than 10 seconds) | 0 |
| | | Moderate (limb upheld less than 10 seconds) | 1 |
| | | Severe (patient unable to raise arm against gravity) | 2 |
| Leg Motor Function | Extending the leg of the patient 30° (in supine) | Normal to Mild (limb upheld more than 5 seconds) | 0 |
| | | Moderate (limb upheld less than 5 seconds) | 1 |
| | | Severe (patient unable to raise leg against gravity) | 2 |
| Head & Gaze Deviation | Observe eyes and head deviation to one side | Absent (eye movements to both sides were possible and no head deviation was observed) | 0 |
| | | Present (eyes and head deviation to one side was observed) | 1 |
| Aphasia (R side) | Difficulty understanding spoken or written words. Ask patient to follow two simple commands: 1. Close your eyes. 2. Make a fist. | Normal (performs both tasks requested correctly) | 0 |
| | | Moderate (performs only 1 of 2 tasks requested correctly) | 1 |
| | | Severe (Cannot perform either task requested correctly) | 2 |
| Agnosia (L side) | Inability to recognize familiar objects. Ask patient: 1. "Whose arm is this?" (while showing the affected arm) 2. "Can you move your arm?" | Normal (recognizes arm, and attempts to move arm) | 0 |
| | | Moderate (does not recognize arm or is unaware of arm) | 1 |
| | | Severe (does not recognize arm and is unaware of arm) | 2 |
| If RACE >5, consider Comprehensive Stroke Center | | RACE SCALE TOTAL | |

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Time of Onset or Last Seen Normal:** Interview patient, family members, and other witnesses to determine Last Known Well (LKW) time and time of Symptom Discovery.
- **Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT "about 45 minutes ago.") Without this information patient may not be able to receive thrombolytics at facility.**
- **For patients with "Woke up and noticed stroke," Time starts when patient went to sleep or was last awake and was last known normal.**
- **Attempt to identify possible stroke mimics (eg. Seizure, migraine, intoxication) and determine if patient has pre-existing substantial disability (need for nursing home care or inability to walk without help from others).**
- **Encourage family to go directly to Emergency Department if not transported with patient and obtain mobile number of next of kin and witnesses**
- **Patients who are eligible for IV Alteplase if transported to nearest Acute Stroke Ready Hospital (ASRH) or PSC should not be rerouted to a CSC or Endovascular Treatment-capable Center if doing so would result in delay that would make them ineligible for IV Alteplase**
- **Air Medical:** Important for EMS to be aware of role of air medical. May be needed to transfer a stroke patient to a geographically distant hospital that is capable of providing an advanced level of stroke care.
- **With a duration of symptoms of less than 3.5 HOURS or UNDETERMINED, scene times should be limited to ≤ 15 minutes and the patient should be transported to capable stroke receiving facility. In-field notification of receiving facility should be performed and transport times should be minimized.**
- **Collect a list of current medications (especially anticoagulants) and obtain patient history including co-morbid conditions (eg. Serious kidney or liver disease, recent surgery, procedures or stroke) that may impact treatment decisions.**