STROKES CAN HAPPEN AT ANY AGE Pediatric stroke can happen in infants, children and even before birth.

PERINATAL STROKE

Last few months of pregnancy to 1-month-old

CHILDHOOD STROKE

1-month-old to 18 years

RISK FACTORS

The cause in most perinatal strokes remains unknown.

Risk factors that could lead to stroke include:

- · Congenital heart disease
- · Disorders of the placenta
- · Blood clotting disorders
- Infections (e.g. Meningitis)

Risk factors in children ≠ Risk factors in older adults Risk factors for children include:



- Congenital heart disease
- Diseases affecting the brain's arteries
- Infections affecting the brain or other organs
- Head trauma
- Sickle cell disease
- Autoimmune disorders

No previous risk factor is identified in about half of childhood stroke cases.

WARNING SIGNS

Signs of a perinatal stroke may go unrecognized for months or years because the signs can be subtle.

NEWBORNS:

Seizures may be an early sign:

- · Repetitive twitching of face, arm or leg
- · Apnea (pauses in breathing) associated with staring

DEVELOPING CHILDREN:

- · Decreased movement or weakness on one side of the body
- Showing a hand preference, or consistently reaching out with only one hand before 1 year of age

Signs are often missed in children because there is a lack of awareness that strokes can happen in this age group.

FACE

STROKE SIGNS ▶

ADDITIONAL SIGNS IN CHILDREN INCLUDE:

DROOPING WFAKNESS

SPEECH

TIME TO

DIFFICULTY

• Severe sudden headache, especially with vomiting and sleepiness

- · Weakness or numbness on one side of the body difficulty speaking or understanding others
- · Vision loss or double vision
- Severe dizziness or loss of coordination
- New-onset of seizures usually on one side of the body

TIME IS BRAIN AT ANY AGE

NEWBORNS:

Quick recognition > Prompt medical evaluation and treatment

BABIES:

Early diagnosis
Rehabilitation treatment can start while a young brain is still developing



DON'T DELAY!

Prompt diagnosis and treatment of stroke in children is as critical as it is in adults.



